

GASTROPARESIS DIET FOR DELAYED STOMACH EMPTYING

MIDWEST GASTROENTEROLOGY

(816) 836-2200

Purpose

Gastroparesis is the medical term for delayed stomach emptying. During the process of digestion, the stomach must contract to empty itself of food and liquid. Normally, it contracts about three times a minute. This empties the stomach within 90-120 minutes after eating. If contractions are sluggish or less frequent, stomach emptying is delayed. This results in bothersome and sometimes serious symptoms, as well as malnutrition, because food is not being digested properly.

Gastroparesis may be caused by various conditions such as diabetes mellitus, certain disorders of the nervous system, or certain drugs. Often however, no cause can be found. Usually, the physician prescribes medication to stimulate the stomach to contract. The purpose of the gastroparesis diet is to reduce symptoms and maintain adequate fluids and nutrition. There are three steps to the diet:

STEP 1 DIET consists of liquids, which usually leave the stomach quickly by gravity alone. Liquids prevent dehydration and keep the body supplied with vital salts and minerals.

STEP 2 DIET provides additional calories by adding a small amount of dietary fat - - less than 40 gm. Each day. For patients with gastroparesis, fatty foods and oils should be restricted, because they delay stomach emptying. However, patients at the Step 2 level are usually able to tolerate this amount.

STEP 3 DIET is designed for long-term maintenance. Fat is limited to 50 gm. per day, and fibrous foods are restricted, because many plant fibers cannot be digested.

Nutrition Facts

The STEP 1 Gastroparesis Diet is inadequate in all nutrients except sodium and potassium. It should not be continued for more than three days without additional nutritional support. STEP 2 and STEP 3 Gastroparesis Diets may be inadequate in Vitamins A and C, and the mineral iron. A multi-vitamin supplement is usually prescribed.

SPECIAL CONSIDERATIONS

1. Diets must be tailored to the individual patient. This is because the degree of gastroparesis may range from severe and long-standing to mild and easily corrected. Patients may also have various medical conditions to be considered. For example, diabetes patients with gastroparesis are allowed sugar-containing liquids on the STEP 2 diet, because that is their only source of carbohydrate. On the STEP 2 and STEP 3 diets, these patients should avoid concentrated sweets. These are noted with an asterisk (*) on the food lists.
2. On all of the diets, liquids and food should be eaten in small, frequent meals. This helps maintain nutrition.

STEP 1 FOOD GROUPS

Group	Recommended	Avoid
Milk & milk products	None	All
Vegetables	None	All
Fruits	None	All
Breads & grain	Plain saltine crackers	All others
Meats & meat substitutes	None	All
Fats & oils	None	All
Beverages	Gatorade and soft drinks (sipped slowly throughout the day)	All others
Soups	Fat-free consommé' and bouillon	All others

SAMPLE MENU STEP 1

Breakfast	Lunch	Dinner
<ul style="list-style-type: none"> • Gatorade ½ cup • Ginger ale ½ cup • Bouillon ¾ cup • Saltine crackers 6 	<ul style="list-style-type: none"> • Gatorade ½ cup • Coke ½ cup • Bouillon ¾ cup • Saltine crackers 6 	<ul style="list-style-type: none"> • Gatorade ½ cup • Sprite ½ cup • Bouillon ¾ cup • Saltine crackers 6

THIS SAMPLE DIET PROVIDES THE FOLLOWING:

Calories	790	Fat	9 gm.
Protein	11 gm.	Sodium	3531 mg.
Carbohydrates	156 gm.	Potassium	244 mg.

STEP 2 FOOD GROUPS

Group	Recommended	Avoid
Milk & milk products	Skim milk, products made with skim milk, low-fat yogurt, low-fat cheeses	Whole milk products, creams (sour, light, heavy, whipping), half & half
Soups	Fat-free consommé' & bouillon, soups made from skim milk, & fat-free broths containing pasta or noodles and allowed vegetables	Soups made with cream, whole milk, or broths containing fat
Bread & grains	Breads & cereals, cream of wheat, pasta, white rice, egg noodles, low-fat crackers	Oatmeal; whole grain rice, cereal, bread
Meat & meat substitutes	Eggs, peanut butter (maximum 2 Tbs/day)	Beef; poultry; fish; pork products; dried beans, peas, & lentils
Vegetables	Vegetable juice (tomato, V-8); well-cooked vegetables w/o skins (acorn squash, beets, carrots, mushrooms, potatoes, spinach, summer squash, yams)	All raw vegetables; cooked vegetables w/skins; beans (green, wax, lima), broccoli, Brussels sprouts, cabbage, cauliflower, celery, corn, eggplant, onions, peas, peppers, pea pods, sauerkraut, turnips, water chestnuts, zucchini
Fruits	Apple, cranberry, grape, pineapple, & prune juices; canned fruits w/o skins	Citrus juices, all fresh and dried fruits, canned fruits w/skins (apricots,

Fruits (continued)	(applesauce, peaches, pears); strained tomato sauce	cherries, blueberries, fruit cocktail, oranges, grapefruit, pineapple, plums, persimmons)
Fats & oils	Any type of fat, but only in small amounts	None
Sweets & desserts	Hard candies, caramels, puddings & custards made from skim milk, frozen yogurt, fruit ice, gelatin, ice milk, jelly, honey, syrups	High-fat desserts (cakes, pies, cookies, pastries, ice cream), fruit preserves
Beverages	Gatorade*, soft drinks*, (sipped slowly throughout the day)	All others, except allowed juices
*Concentrated sweets		

SAMPLE MENU STEP 2

Breakfast	Lunch	Dinner
<ul style="list-style-type: none"> • Skim milk $\frac{1}{2}$ cup • Poached egg 1 • White toast slice • Apple juice $\frac{1}{2}$ cup 	<ul style="list-style-type: none"> • Mozzarella cheese 2 oz. • Saltine crackers 6 • Chicken noodle soup $\frac{3}{4}$ cup • Gatorade $\frac{1}{2}$ cup 	<ul style="list-style-type: none"> • Peanut butter 1 Tbs. • Saltine crackers 6 • Vanilla pudding $\frac{1}{2}$ cup • Grape juice $\frac{1}{2}$ cup
Morning Snack	Afternoon Snack	Evening Snack
<ul style="list-style-type: none"> • Ginger ale $\frac{1}{2}$ cup • Canned pears $\frac{1}{2}$ cup 	<ul style="list-style-type: none"> • Skim milk $\frac{1}{2}$ cup • Cornflakes $\frac{1}{2}$ cup • Sugar 2 tsp. 	<ul style="list-style-type: none"> • Frozen yogurt $\frac{1}{2}$ cup • Saltine crackers 6

THIS SAMPLE DIET PROVIDES THE FOLLOWING:

Calories	1343	Fat	35 gm.
Protein	52 gm.	Sodium	2639 mg.
Carbohydrates	206 gm.	Potassium	1411 mg.

STEP 3 FOODS GROUPS

Group	Recommended	Avoid
Milk & milk products	Skim milk, products made with skim milk, low-fat yogurt, low-fat cheeses	Whole milk products, creams (sour, light, heavy, whipping), half & half
Soups	Fat-free consommé' & bouillon, soups made from skim milk, & fat- free broths containing pasta or noodles and allowed vegetables	Soups made with cream, whole milk, or broths containing fat
Fruits	Fruit juices, canned fruits w/o skins (applesauce, peaches, pears)	All fresh & dried fruits, canned fruits w/skins (apricots, cherries, plums, blueberries, fruit cocktail, oranges, grapefruit, pineapple, persimmons)
Meat & meat substitutes	Eggs, peanut butter (2 Tbs./day), poultry, fish, lean ground beef	Fibrous meats (steaks, roasts, chops), dried beans, peas, lentils
Fats & oils	Any type of fat, but only in small amounts	None
Breads & grains	Breads & cereals, cream of wheat, pasta, white rice, egg noodles, low- fat crackers	Oatmeal; whole grain rice, cereal, bread

Vegetables	Vegetables juices (tomato V-8), well-cooked vegetables w/o skins (acorn squash, beets, carrots, mushrooms, potatoes, spinach, summers squash, strained tomato sauce, yams)	All raw vegetables; cooked vegetables w/skins; beans (green, wax, lima), broccoli, Brussels sprouts, cabbage, cauliflower, celery, corn, eggplant, onions, peas, peppers, pea pods, sauerkraut, turnips, water chestnuts, zucchini
Sweets & desserts	Hard candies, caramels, puddings & custards made from skim milk, frozen yogurt, fruit ice, gelatin, ice milk, jelly, honey, syrups	High-fat desserts (cakes, pies, cookies, pastries, ice cream), fruit preserves
Beverages	Gatorade*, soft drinks* (sipped slowly throughout the day), coffee, tea, water (note: non-caloric beverages should be limited if patient cannot maintain adequate caloric intake)	All others, except allowed juices
*Concentrated Sweets		

SAMPLE MENU STEP 3

Breakfast	Lunch	Dinner
<ul style="list-style-type: none"> • Skim milk $\frac{1}{2}$ cup • Cream of wheat $\frac{1}{2}$ cup • Sugar 2 tsp. • Orange juice $\frac{1}{2}$ cup • White toast 1 slice • Margarine 1 tsp. • Jelly 1 tbs. 	<ul style="list-style-type: none"> • Tuna fish 2 oz. • Low-fat mayonnaise 2 tbs. • White bread 2 slices • Canned peaches $\frac{1}{2}$ cup • Gatorade 1.2 cup 	<ul style="list-style-type: none"> • Baked chicken 2 oz. • White rice $\frac{1}{2}$ cup • Cooked beets $\frac{1}{2}$ cup • Dinner roll 1 • Skim milk $\frac{1}{2}$ cup • Margarine 2 tsp.
Morning Snack	Afternoon Snack	Evening Snack
<ul style="list-style-type: none"> • Low-fat yogurt $\frac{1}{2}$ cup • Sprite $\frac{1}{2}$ cup 	<ul style="list-style-type: none"> • Chocolate pudding $\frac{1}{2}$ cup • Ginger ale $\frac{1}{2}$ cup 	<ul style="list-style-type: none"> • Ice milk $\frac{1}{2}$ cup • Pretzels 2

THIS SAMPLE DIET PROVIDES THE FOLLOWING:

Calories	1822	Fat	42 gm.
Protein	75 gm.	Sodium	2234 mg.
Carbohydrates	286 gm.	Potassium	2467 mg.